STIC Search Report
EIC 1700

STIC Database Tracking Number: 201572

TO: Ben Sackey
Location: REM 5B31
Art Unit : 1626
September 14, 2006

Case Serial Number: 10/735029

From: Kathleen Fuller
Location: EIC 1700
REMSEN 4B28
Phone: 571/272-2505
Kathleen.Fuller@uspto.gov

Search Notes

```
=> FILE REG
FILE 'REGISTRY' ENTERED AT 10:19:04 ON 14 SEP 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file
provided by InfoChem.

STRUCTURE FILE UPDATES:   13 SEP 2006  HIGHEST RN 906624-07-5
DICTIONARY FILE UPDATES:  13 SEP 2006  HIGHEST RN 906624-07-5

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 30, 2006

  Please note that search-term pricing does apply when
  conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and
predicted properties as well as tags indicating availability of
experimental property data in the original document.  For information
on property searching in REGISTRY, refer to:

http://www.cas.org/ONLINE/UG/regprops.html

=> FILE HCAPL
FILE 'HCAPLUS' ENTERED AT 10:19:07 ON 14 SEP 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)


Copyright of the articles to which records in this database refer is
held by the publishers listed in the PUBLISHER (PB) field (available
for records published or updated in Chemical Abstracts after December
26, 1996), unless otherwise indicated in the original publications.
The CA Lexicon is the copyrighted intellectual property of the
the American Chemical Society and is provided to assist you in searching
databases on STN.  Any dissemination, distribution, copying, or storing
of this information, without the prior written consent of CAS, is
strictly prohibited.

FILE COVERS 1907 - 14 Sep 2006  VOL 145 ISS 12
FILE LAST UPDATED: 13 Sep 2006  (20060913/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

 This file contains CAS Registry Numbers for easy and accurate
 substance identification.

=> D QUE
L2             5 SEA FILE=REGISTRY ABB=ON  (109-07-9/BI OR 112811-59-3/BI OR
                 112811-72-0/BI OR 121577-32-0/BI OR 713503-65-2/BI)
L4             1 SEA FILE=REGISTRY ABB=ON  L2 AND GATIFLO?
L5          1086 SEA FILE=HCAPLUS ABB=ON  L4
L6            10 SEA FILE=HCAPLUS ABB=ON  L5(L)CRYSTAL?
L8            16 SEA FILE=HCAPLUS ABB=ON  L5 AND CRYSTAL?(4A)GATIFLOXACIN?
L10            3 SEA FILE=REGISTRY ABB=ON  L2 AND 4/NR
L11            2 SEA FILE=REGISTRY ABB=ON  L10 NOT L4
```

```
L12           6 SEA FILE=HCAPLUS ABB=ON  L11
L13           1 SEA FILE=HCAPLUS ABB=ON  L12 AND CRYSTAL?
L14          16 SEA FILE=HCAPLUS ABB=ON  L8 OR L6 OR L13


=> D L14 BIB ABS IND HITSTR 1-16

L14  ANSWER 1 OF 16  HCAPLUS  COPYRIGHT 2006 ACS on STN
AN   2005:1311539  HCAPLUS
DN   144:22824
TI   Novel crystalline forms of gatifloxacin
IN   Satyanarayana, Chava; Ramanjaneyulu, Gorantla Seeta; Kumar, Indukuri
     Venkata Sunil
PA   Matrix Laboratories Ltd, India
SO   PCT Int. Appl., 23 pp.
     CODEN: PIXXD2
DT   Patent
LA   English
FAN.CNT 1
     PATENT NO.          KIND   DATE        APPLICATION NO.         DATE
     --------------      ----   --------    --------------------    --------
PI   WO 2005118546        A1    20051215    WO 2005-IN166           20050525
         W:  AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH,
             CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,
             GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ,
             LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA,
             NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK,
             SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU,
             ZA, ZM, ZW
         RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,
             AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,
             EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT,
             RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML,
             MR, NE, SN, TD, TG
PRAI IN 2004-CH521         A     20040604
     IN 2004-CH522         A     20040604
     IN 2004-CH523         A     20040604
AB   A process for purification of polymorphic form of Gatifloxacin is presented.
     Gatifloxacin is dissolved about 15 - 50 vols. of methanol, thereby
     removing insolubles, followed by adding organic base to the solution,. By
     maintaining the solution at temperature of 30 °C to 70 °C, for about
     20 min to 4 h, followed by gradual cooling and maintaining the reaction
     mass to -10 to 20 °C for about 1 - 4 h, isolation and drying at
     temperature of about 45 °C to 65 °C enables gatifloxacin
     to crystallized in three novel polymorphs.
IC   ICM  C07D215-56
CC   27-17 (Heterocyclic Compounds (One Hetero Atom))
     Section cross-reference(s): 75
ST   crystallog purifn gatifloxacin polymorph;
     crystal mol structure gatifloxacin polymorph
IT   Crystal structure
     Molecular structure
        (of three unique polymorphs of gatifloxacin)
IT   Conformation
     Conformers
     Differential scanning calorimetry
     Polymorphism (crystal)
     Purification
     X-ray diffraction
        (process for the crystallog. purification of three unique polymorphs of
```

gatifloxacin)
IT 67-56-1, Methanol, uses
RL: NUU (Other use, unclassified); USES (Uses)
(process for the crystallog. purification of three unique polymorphs of gatifloxacin)
IT **112811-59-3P, Gatifloxacin**
RL: PRP (Properties); PUR (Purification or recovery); PREP (Preparation)
(process for the **crystallog.** purification of three unique polymorphs of gatifloxacin)
IT **112811-59-3P, Gatifloxacin**
RL: PRP (Properties); PUR (Purification or recovery); PREP (Preparation)
(process for the **crystallog.** purification of three unique polymorphs of gatifloxacin)
RN 112811-59-3 HCAPLUS
CN 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo- (9CI) (CA INDEX NAME)



RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L14 ANSWER 2 OF 16 HCAPLUS COPYRIGHT 2006 ACS on STN
AN 2005:572069 HCAPLUS
DN 143:172897
TI Preparation of Gatifloxacin hydrobromide and application as antibacterial agent
IN Tang, Xiaodong; Tang, Xudong
PA Hainan Kangliyuan Pharmaceutical Industry Co., Ltd., Peop. Rep. China
SO Faming Zhuanli Shenqing Gongkai Shuomingshu, No pp. given
CODEN: CNXXEV
DT Patent
LA Chinese
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	CN 1548435	A	20041124	CN 2003-116874	20030509
PRAI	CN 2003-116874		20030509		

GI



I

AB The present invention discloses preparation of Gatifloxacin hydrobromide (I) and its application. The preparation process adopts Gatifloxacin as main material, and includes reaction with hydrobromic acid in organic solvent or water at 40-50° C for 1-5 h to produce the salt, and subsequent **crystallization** to obtain **Gatifloxacin** hydrobromide product. The compound has high water solubility, high stability and less irritation, and its Gatifloxacin component produces antibacterial effect in human body to treat various infectious diseases.

IC ICM C07D401-04
ICS A61K031-496; A61P031-04

CC 28-17 (Heterocyclic Compounds (More Than One Hetero Atom))
Section cross-reference(s): 10

ST Gatifloxacin hydrobromide prepn antibacterial agent

IT Infection
(bacterial; preparation of the hydrobromide salt of antibacterial agent Gatifloxacin)

IT Antibacterial agents
Human
(preparation of the hydrobromide salt of antibacterial agent Gatifloxacin)

IT **112811-59-3**, Gatifloxacin
RL: PAC (Pharmacological activity); RCT (Reactant); THU (Therapeutic use); BIOL (Biological study); RACT (Reactant or reagent); USES (Uses)
(preparation of the hydrobromide salt of antibacterial agent Gatifloxacin)

IT 316819-22-4P
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of the hydrobromide salt of antibacterial agent Gatifloxacin)

IT **112811-59-3**, Gatifloxacin
RL: PAC (Pharmacological activity); RCT (Reactant); THU (Therapeutic use); BIOL (Biological study); RACT (Reactant or reagent); USES (Uses)
(preparation of the hydrobromide salt of antibacterial agent Gatifloxacin)

RN 112811-59-3 HCAPLUS

CN 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo- (9CI) (CA INDEX NAME)



L14 ANSWER 3 OF 16 HCAPLUS COPYRIGHT 2006 ACS on STN
AN 2005:451362 HCAPLUS
DN 142:487546
TI Preparation of a **crystalline** form of **gatifloxacin** which has a stable water content
IN Cosme Gomez, Antonio; Villasante Prieto, Javier; Palomo Nicolau, Francisco Eugenio
PA Quimica Sintetica, S. A., Spain
SO PCT Int. Appl., 19 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 2005047262	A1	20050526	WO 2004-IB3652	20041105
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
ES 2232311	A1	20050516	ES 2003-2643	20031113
ES 2232311	B1	20060801		
PRAI ES 2003-2643	A	20031113		

AB A **crystalline** form of **gatifloxacin**, obtained by a process that comprises recrystn. of crude gatifloxacin in methanol, which is stable with a water content ranging between 2.5-4.5%, is prepared and is claimed for use in the preparation of pharmaceutical formulations for the treatment of bacterial infections.
IC ICM C07D215-56
ICS A61K031-496
CC 63-6 (Pharmaceuticals)
Section cross-reference(s): 28, 75
ST **gatifloxacin cryst** form prepn stable water content; polymorphic **gatifloxacin cryst** form prepn stable water content
IT Infection
(bacterial; preparation of a **crystalline** form of **gatifloxacin** which has a stable water content for use in pharmaceutical formulations for the treatment of)
IT Cooling
Crystallization
Filtration
Heating
(in the preparation of a **crystalline** form of **gatifloxacin** which has a stable water content)
IT Polymorphism (crystal)
(preparation of a **crystalline** form of **gatifloxacin** which has a stable water content)
IT Drug delivery systems
(preparation of a **crystalline** form of **gatifloxacin** which has a stable water content for use in)

IT Antibiotics
(preparation of a **crystalline** form of **gatifloxacin** which has a stable water content for use in pharmaceutical formulations as)
IT Drying
(vacuum; in the preparation of a **crystalline** form of **gatifloxacin** which has a stable water content)
IT **112811-59-3, Gatifloxacin**
RL: PEP (Physical, engineering or chemical process); PRP (Properties); PYP (Physical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)
(preparation of a **crystalline** form of **gatifloxacin** which has a stable water content)
IT 67-56-1, Methanol, uses
RL: NUU (Other use, unclassified); USES (Uses)
(solvent; preparation of a **crystalline** form of **gatifloxacin** which has a stable water content)
IT **112811-59-3, Gatifloxacin**
RL: PEP (Physical, engineering or chemical process); PRP (Properties); PYP (Physical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)
(preparation of a **crystalline** form of **gatifloxacin** which has a stable water content)
RN 112811-59-3 HCAPLUS
CN 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo- (9CI) (CA INDEX NAME)



RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L14 ANSWER 4 OF 16 HCAPLUS COPYRIGHT 2006 ACS on STN
AN 2005:451361 HCAPLUS
DN 142:487545
TI Process for the preparation of a non-hygroscopic polymorphic **crystalline** form of **gatifloxacin**
IN Cosme Gomez, Antonio; Villasante Prieto, Javier; Palomo Nicolau, Francisco Eugenio
PA Quimica Sintetica, S. A., Spain
SO PCT Int. Appl., 24 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 2005047261	A1	20050526	WO 2004-IB3600	20041105

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,

LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI,
NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY,
TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,
AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO,
SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG

ES 2232310 A1 20050516 ES 2003-2642 20031113
ES 2232310 B1 20060801
PRAI ES 2003-2642 A 20031113

AB A non-hygroscopic polymorphic (I) **crystalline** form of **gatifloxacin**, which contains 8-9% water, which is characterized by a specific powder X-ray diffraction pattern, is prepared by heating crude I in methanol and cooling it with stirring to crystallization I is claimed for its use as an active ingredient in the preparation of antibiotic pharmaceutical formulations.

IC ICM C07D215-56
ICS A61K031-496; A61P031-00

CC 63-6 (Pharmaceuticals)
Section cross-reference(s): 28, 75

ST **gatifloxacin** nonhygroscopic **crystal** polymorph prepn; antibiotic **gatifloxacin** nonhygroscopic **crystal** polymorph prepn

IT Infection
(bacterial; process for the preparation of a non-hygroscopic polymorphic **crystalline** form of **gatifloxacin** for use in the treatment of)

IT Cooling
Crystallization
Filtration
Heating
(in a process for the preparation of a non-hygroscopic polymorphic **crystalline** form of **gatifloxacin**)

IT Antibiotics
Polymorphism (crystal)
(process for the preparation of a non-hygroscopic polymorphic **cryst** . form of **gatifloxacin**)

IT Drug delivery systems
(process for the preparation of a non-hygroscopic polymorphic **cryst** . form of **gatifloxacin** for use in)

IT **112811-59-3**, Gatifloxacin
RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(process for the preparation of a non-hygroscopic polymorphic **cryst** . form of **gatifloxacin**)

IT 67-56-1, Methanol, uses
RL: NUU (Other use, unclassified); USES (Uses)
(solvent; process for the preparation of a non-hygroscopic polymorphic **crystalline** form of **gatifloxacin**)

IT **112811-59-3**, Gatifloxacin
RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(process for the preparation of a non-hygroscopic polymorphic **cryst** . form of **gatifloxacin**)

RN 112811-59-3 HCAPLUS
CN 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo- (9CI) (CA INDEX NAME)



RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L14 ANSWER 5 OF 16 HCAPLUS COPYRIGHT 2006 ACS on STN
AN 2005:220275 HCAPLUS
DN 143:240624
TI The **crystal** structure of a **Gatifloxacin** complex and its fluorescent property
AU Li, Yong-Hua; Tang, Yun-Zhi; Huang, Xue-Feng; Xiong, Ren-Gen
CS Coordination Chemistry Institute, The State Key Laboratory of Coordination Chemistry, Nanjing University, Nanjing, Peop. Rep. China
SO Zeitschrift fuer Anorganische und Allgemeine Chemie (2005), 631(4), 639-641
CODEN: ZAACAB; ISSN: 0044-2313
PB Wiley-VCH Verlag GmbH & Co. KGaA
DT Journal
LA English
OS CASREACT 143:240624
AB The hydrothermal reaction of Cd(ClO4)2·6H2O and Gatifloxacin (Gati) affords a binuclear complex of Gatifloxacin, [Cd(Gati)(py)3]2(ClO4)4 (1), which was characterized by x-ray crystallog. anal., IR and fluorescent spectroscopy. Crystal data for 1, C34H36CdCl2FN6O12: monoclinic, space group P21/c, a 18.3642(5), b 12.8156(4), c 16.7210(5) Å, β 97.931(1)°, Z = 4. The local surrounding of the CdII ion is a slightly distorted octahedron. Solid-state fluorescence measurements of 1 at room temperature show an emission peak at 439 nm.
CC 78-7 (Inorganic Chemicals and Reactions)
Section cross-reference(s): 73, 75
ST cadmium Gatifloxacin dinuclear complex prepn structure fluorescence; **crystal** structure cadmium **Gatifloxacin** dinuclear complex
IT **Crystal** structure
Fluorescence
Molecular structure
(of cadmium **Gatifloxacin** dinuclear complex)
IT 862771-77-5P
RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)
(preparation and crystal structure and fluorescence of)
IT **112811-59-3**, Gatifloxacin
RL: RCT (Reactant); RACT (Reactant or reagent)
(reactant for preparation of cadmium Gatifloxacin dinuclear complex)
IT **112811-59-3**, Gatifloxacin
RL: RCT (Reactant); RACT (Reactant or reagent)
(reactant for preparation of cadmium Gatifloxacin dinuclear complex)
RN 112811-59-3 HCAPLUS
CN 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo- (9CI) (CA INDEX NAME)



RE.CNT 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L14 ANSWER 6 OF 16 HCAPLUS COPYRIGHT 2006 ACS on STN
AN 2004:1016034 HCAPLUS
DN 142:11555
TI Purification methods for gatifloxacin and preparation of a polymorphic **crystalline** form of **gatifloxacin** hemihydrate
IN Reddy, Bandi Parthasaradhi; Reddy, Kura Rathnakar; Reddy, Rapolu Raji; Reddy, Dasari Muralidhara; Reddy, Jonnala Sambi
PA Hetero Drugs Limited, India
SO PCT Int. Appl., 18 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004101547	A1	20041125	WO 2003-IN191	20030519
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	AU 2003304118	A1	20041203	AU 2003-304118	20030519
PRAI	WO 2003-IN191	A	20030519		

AB A process for the purification of gatifloxacin contaminated with impurities comprises: (A) adjusting the pH of the suspension of impure gatifloxacin in water to <4.5 until the solids are in solution; (B) separating the impurities by solvent extraction and/or adsorption; (C) readjusting the pH to 6-8 with a base; and (D) isolating purified **crystalline gatifloxacin.** A stable novel **crystalline** form of **gatifloxacin** hemihydrate is also presented.

IC ICM C07D401-04
ICS C07D215-56; A61K031-496
CC 63-6 (Pharmaceuticals)
Section cross-reference(s): 28, 75
ST **gatifloxacin** hemihydrate prepn **crystal** polymorphism; purifn gatifloxacin
IT Alkali metal hydroxides
Alkaline earth hydroxides
Amines, reactions
Carbonates, reactions

Phosphates, reactions
RL: RGT (Reagent); RACT (Reactant or reagent)
(bases; in purification methods for gatifloxacin and preparation of a polymorphic
crystalline form of **gatifloxacin** hemihydrate)

IT Hydrocarbons, uses
RL: NUU (Other use, unclassified); USES (Uses)
(chloro, solvents; in purification methods for gatifloxacin and preparation of a
polymorphic **crystalline** form of **gatifloxacin** hemihydrate)

IT Amines, reactions
RL: RGT (Reagent); RACT (Reactant or reagent)
(heterocyclic, bases; in purification methods for gatifloxacin and preparation of
a polymorphic **crystalline** form of **gatifloxacin** hemihydrate)

IT Adsorption
Extraction
(in purification methods for gatifloxacin and preparation of a polymorphic **crystalline** form of **gatifloxacin** hemihydrate)

IT Bases, reactions
RL: RGT (Reagent); RACT (Reactant or reagent)
(in purification methods for gatifloxacin and preparation of a polymorphic **crystalline** form of **gatifloxacin** hemihydrate)

IT Acids, reactions
RL: RGT (Reagent); RACT (Reactant or reagent)
(inorg.; in purification methods for gatifloxacin and preparation of a polymorphic **crystalline** form of **gatifloxacin** hemihydrate)

IT Acids, reactions
RL: RGT (Reagent); RACT (Reactant or reagent)
(organic; in purification methods for gatifloxacin and preparation of a polymorphic
crystalline form of **gatifloxacin** hemihydrate)

IT Polymorphism (**crystal**)
(purification methods for **gatifloxacin** and preparation of a polymorphic **crystalline** form of **gatifloxacin** hemihydrate)

IT Drug delivery systems
(purification methods for gatifloxacin and preparation of a polymorphic **crystalline** form of **gatifloxacin** hemihydrate for use in)

IT Esters, uses
Ethers, uses
Hydrocarbons, uses
Ketones, uses
RL: NUU (Other use, unclassified); USES (Uses)
(solvents; in purification methods for gatifloxacin and preparation of a polymorphic **crystalline** form of **gatifloxacin** hemihydrate)

IT 7440-44-0, Activated carbon, uses
RL: NUU (Other use, unclassified); PEP (Physical, engineering or chemical process); PYP (Physical process); PROC (Process); USES (Uses)
(activated; in purification methods for gatifloxacin and preparation of a polymorphic **crystalline** form of **gatifloxacin** hemihydrate)

IT 75-50-3, Trimethylamine, reactions 109-02-4, N-Methylmorpholine 109-73-9, Butylamine, reactions 109-89-7, Diethylamine, reactions 110-89-4, Piperidine, reactions 141-43-5, Ethanolamine, reactions 497-19-8, Sodium carbonate, reactions 1310-58-3, Potassium hydroxide, reactions 1310-73-2, Sodium hydroxide, reactions 7601-54-9, Trisodium

phosphate 7664-41-7, Ammonia, reactions
RL: RGT (Reagent); RACT (Reactant or reagent)
(base; in purification methods for gatifloxacin and preparation of a polymorphic
crystalline form of **gatifloxacin** hemihydrate)

IT 64-18-6, Formic acid, reactions 64-19-7, Acetic acid, reactions 76-05-1, Trifluoroacetic acid, reactions 7647-01-0, Hydrochloric acid, reactions 7664-38-2, Phosphoric acid, reactions 7664-93-9, Sulfuric acid, reactions
RL: RGT (Reagent); RACT (Reactant or reagent)
(in purification methods for gatifloxacin and preparation of a polymorphic **crystalline** form of **gatifloxacin** hemihydrate)

IT 404858-36-2P, Gatifloxacin hemihydrate
RL: PEP (Physical, engineering or chemical process); PRP (Properties); PYP (Physical process); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)
(purification methods for gatifloxacin and preparation of a polymorphic **crystalline** form of **gatifloxacin** hemihydrate)

IT **112811-59-3P**, Gatifloxacin
RL: PEP (Physical, engineering or chemical process); PUR (Purification or recovery); PYP (Physical process); RCT (Reactant); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); RACT (Reactant or reagent); USES (Uses)
(purification methods for gatifloxacin and preparation of a polymorphic **crystalline** form of **gatifloxacin** hemihydrate)

IT 67-66-3, Chloroform, uses 75-09-2, Dichloromethane, uses 108-10-1, MIBK 108-20-3, Isopropyl ether 108-21-4, Isopropyl acetate 108-88-3, Toluene, uses 110-19-0, Isobutyl acetate 141-78-6, Ethyl acetate, uses
RL: NUU (Other use, unclassified); USES (Uses)
(solvent; in purification methods for gatifloxacin and preparation of a polymorphic **crystalline** form of **gatifloxacin** hemihydrate)

IT 7732-18-5, Water, reactions
RL: NUU (Other use, unclassified); RCT (Reactant); RACT (Reactant or reagent); USES (Uses)
(solvent; in purification methods for gatifloxacin and preparation of a polymorphic **crystalline** form of **gatifloxacin** hemihydrate)

IT **112811-59-3P**, Gatifloxacin
RL: PEP (Physical, engineering or chemical process); PUR (Purification or recovery); PYP (Physical process); RCT (Reactant); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); RACT (Reactant or reagent); USES (Uses)
(purification methods for gatifloxacin and preparation of a polymorphic **crystalline** form of **gatifloxacin** hemihydrate)

RN 112811-59-3 HCAPLUS

CN 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo- (9CI) (CA INDEX NAME)



RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L14 ANSWER 7 OF 16 HCAPLUS COPYRIGHT 2006 ACS on STN
AN 2004:857586 HCAPLUS
DN 141:320019
TI Preparation of novel **crystalline** forms of **gatifloxacin**
IN Parthasaradhi, Reddy Bandi; Rathnakar, Reddy Kura; Raji, Reddy Rapolu; Muralidhara, Reddy Dasari; Ravikanth, Reddy Meghi
PA Hetero Drugs Limited, India
SO PCT Int. Appl., 19 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 2004087688	A1	20041014	WO 2003-IN135	20030402
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2003230194	A1	20041025	AU 2003-230194	20030402
US 2005085640	A1	20050421	US 2003-510172	20030402
PRAI WO 2003-IN135	A	20030402		

AB The present invention relates to novel **crystalline** forms of **gatifloxacin**, to processes for their preparation and to pharmaceutical compns. containing them. Thus, 1 g of gatifloxacin was mixed with methylene dichloride (50 mL, water content 0.35%), heated to 45° and maintained at this temp for 15 min. The solution formed was cooled to 25° and maintained at 25° and maintained at 25° for 10 h. The separated crystals were filtered to give 0.6 g of gatifloxacin sesquihydrate form H1.
IC ICM C07D401-04
ICS A61K031-495
CC 63-5 (Pharmaceuticals)
ST **cryst gatifloxacin** prepn
IT Crystal morphology
Solvents
(preparation of novel **crystalline** forms of **gatifloxacin**)
IT 56-23-5, Carbon tetrachloride, uses 67-66-3, Chloroform, uses 75-09-2, Methylene dichloride, uses 79-20-9, Methyl acetate 107-06-2, Ethylene dichloride, uses 107-31-3, Methyl formate 108-21-4, Isopropyl acetate 109-94-4, Ethyl formate 123-91-1, 1,4 Dioxane, uses 141-78-6, Ethyl acetate, uses 540-88-5, tert-Butyl acetate
RL: NUU (Other use, unclassified); USES (Uses)
(preparation of novel **crystalline** forms of **gatifloxacin**)
IT **112811-59-3, Gatifloxacin** 180200-66-2, **Gatifloxacin** sesquihydrate
RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(preparation of novel **crystalline** forms of **gatifloxacin**)
IT **112811-59-3, Gatifloxacin**

RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(preparation of novel **crystalline** forms of **gatifloxacin**)
RN 112811-59-3 HCAPLUS
CN 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo- (9CI) (CA INDEX NAME)



L14 ANSWER 8 OF 16 HCAPLUS COPYRIGHT 2006 ACS on STN
AN 2004:531362 HCAPLUS
DN 141:94276
TI **Crystalline** forms of **gatifloxacin**
IN Niddam-Hildesheim, Valerie; Wizel, Shomit; Amir, Ehud; Sterimbaum, Greta
PA Teva Pharmaceutical Industries Ltd., Israel; Teva Pharmaceuticals USA, Inc.
SO PCT Int. Appl., 71 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 3

applicants

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004054583	A1	20040701	WO 2003-US39539	20031212
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	EP 1645274	A1	20060412	EP 2005-77643	20030512
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, SK				
	CA 2510625	AA	20040701	CA 2003-2510625	20031212
	AU 2003300874	A1	20040709	AU 2003-300874	20031212
	US 2004171621	A1	20040902	US 2003-735029	20031212
	EP 1485097	A1	20041215	EP 2003-813395	20031212
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
	JP 2006511618	T2	20060406	JP 2005-508325	20031212
	US 2005288301	A1	20051229	US 2005-208248	20050819
	US 2005288302	A1	20051229	US 2005-208299	20050819
PRAI	US 2002-432961P	P	20021212		
	US 2003-448062P	P	20030215		
	US 2003-465534P	P	20030425		

US 2002-379510P	P	20020510
US 2002-389093P	P	20020614
US 2002-401672P	P	20020806
US 2002-402749P	P	20020812
US 2002-409860P	P	20020910
US 2002-423338P	P	20021101
US 2003-444812P	P	20030203
EP 2003-750112	A3	20030512
US 2003-436736	A3	20030512
WO 2003-US39539	W	20031212

GI



I

AB Provided are novel **crystalline** forms of **gatifloxacin** (I), some of which are DMSO solvates. A DMSO solvate of I was prepared from 1-cyclopropyl-6,7-difluoro-1,4-dihydro-8-methoxy-4-oxo-3-quinolinecarboxylic acid, DMSO, and 2-methylpiperazine. Other **crystal** forms and DMSO solvates were prepared and characterized.

IC ICM A61K031-496
ICS C07D215-56; A61P031-00

CC 63-5 (Pharmaceuticals)
Section cross-reference(s): 28, 75

ST **gatifloxacin crystal** form DMSO solvate

IT **Crystal** morphology
(**crystalline** forms of **gatifloxacin**)

IT **Crystal** structure
(of **gatifloxacin**-DMSO solvate)

IT **112811-59-3, Gatifloxacin 121577-32-0,** 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo-, monohydrochloride
RL: CPS (Chemical process); PEP (Physical, engineering or chemical process); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)
(**crystalline** forms of **gatifloxacin**)

IT **713503-65-2**
RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(**crystalline** forms of **gatifloxacin**)

IT 109-07-9, 2-Methylpiperazine 112811-72-0, 1-Cyclopropyl-6,7-difluoro-1,4-dihydro-8-methoxy-4-oxo-3-quinolinecarboxylic acid
RL: RCT (Reactant); RACT (Reactant or reagent)
(**crystalline** forms of **gatifloxacin**)

IT **112811-59-3, Gatifloxacin 121577-32-0,** 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo-, monohydrochloride
RL: CPS (Chemical process); PEP (Physical, engineering or chemical process); PRP (Properties); THU (Therapeutic use); BIOL (Biological

study); PROC (Process); USES (Uses)
(**crystalline** forms of **gatifloxacin**)

RN 112811-59-3 HCAPLUS
CN 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo- (9CI) (CA INDEX NAME)



RN 121577-32-0 HCAPLUS
CN 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl

IT **713503-65-2**
RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(**crystalline** forms of **gatifloxacin**)

RN 713503-65-2 HCAPLUS
CN 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo-, compd. with sulfinylbis[methane] (9CI) (CA INDEX NAME)

CM 1

CRN 112811-59-3
CMF C19 H22 F N3 O4



CM 2

CRN 67-68-5
CMF C2 H6 O S



RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L14 ANSWER 9 OF 16 HCAPLUS COPYRIGHT 2006 ACS on STN
AN 2003:1006776 HCAPLUS
DN 140:31458
TI Novel **crystalline** forms of **gatifloxacin**
IN Niddam-Hildesheim, Valerie; Wizel, Shlomit; Sterimbaum, Greta; Amir, Ehud
PA Teva Pharmaceuticals Industries Ltd., Israel; Teva Pharmaceuticals USA, Inc.
SO PCT Int. Appl., 25 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 3

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 2003105851	A1	20031224	WO 2003-US19046	20030616
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
EP 1645274	A1	20060412	EP 2005-77643	20030512
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, SK				
CA 2489377	AA	20031224	CA 2003-2489377	20030616
AU 2003243615	A1	20031231	AU 2003-243615	20030616
US 2004038988	A1	20040226	US 2003-462945	20030616
EP 1471911	A1	20041103	EP 2003-760424	20030616
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				

IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK

	JP 2005532364	T2	20051027	JP 2004-512754	20030616
	US 2005288301	A1	20051229	US 2005-208248	20050819
	US 2005288302	A1	20051229	US 2005-208299	20050819
PRAI	US 2002-389093P	P	20020614		
	US 2002-423338P	P	20021101		
	US 2002-379510P	P	20020510		
	US 2002-401672P	P	20020806		
	US 2002-402749P	P	20020812		
	US 2002-409860P	P	20020910		
	US 2002-432961P	P	20021212		
	US 2003-444812P	P	20030203		
	US 2003-448062P	P	20030215		
	EP 2003-750112	A3	20030512		
	US 2003-436736	A3	20030512		
	WO 2003-US19046	W	20030616		

AB Provided are two novel **crystalline** forms of **gatifloxacin** (I), denominated form O and form V, methods for their preparation and pharmaceutical compns. thereof. A method of making the crystalline I comprises the steps of (1) providing, at reflux, a solution of I in acetonitrile, (2) cooling the solution to ambient temperature at a cooling rate of at least 1° per min, whereby a suspension is obtained, (3) further crash cooling the suspension to about 5° or less, (4) isolating the solid from the suspension, and (5) treating the isolated solid with moist gas to obtain form V.

IC ICM A61K031-4725
ICS C07D215-56

CC 63-5 (Pharmaceuticals)

ST **gatifloxacin** polymorph **crystn**

IT **Crystallization**
Differential scanning calorimetry
Polymorphism (**crystal**)
X-ray diffractometry
X-ray reflectivity spectra
(novel **crystalline** forms of **gatifloxacin**)

IT 64-17-5, Ethanol, miscellaneous 75-05-8, Acetonitrile, miscellaneous
RL: MSC (Miscellaneous)
(novel **crystalline** forms of **gatifloxacin**)

IT **112811-59-3, Gatifloxacin** 180200-66-2, **Gatifloxacin** sesquihydrate
RL: PEP (Physical, engineering or chemical process); PRP (Properties); PYP (Physical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)
(novel **crystalline** forms of **gatifloxacin**)

IT **112811-59-3, Gatifloxacin**
RL: PEP (Physical, engineering or chemical process); PRP (Properties); PYP (Physical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)
(novel **crystalline** forms of **gatifloxacin**)

RN 112811-59-3 HCAPLUS

CN 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo- (9CI) (CA INDEX NAME)



RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L14 ANSWER 10 OF 16 HCAPLUS COPYRIGHT 2006 ACS on STN
AN 2003:913003 HCAPLUS
DN 139:386418
TI Novel **crystalline** forms of **gatifloxacin**
IN Niddam-Hildesheim, Valerie; Wizel, Shlomit; Sterimbaum, Greta; Amir, Ehud
PA Teva Pharmaceutical Industries Ltd., Israel; Teva Pharmaceuticals USA, Inc.
SO PCT Int. Appl., 91 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 3

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003094919	A2	20031120	WO 2003-US14811	20030512
	WO 2003094919	A3	20040318		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	AU 2003232113	A1	20031111	AU 2003-232113	20030512
	CA 2485262	AA	20031120	CA 2003-2485262	20030512
	US 2004009989	A1	20040115	US 2003-436736	20030512
	EP 1503762	A2	20050209	EP 2003-750112	20030512
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
	CN 1665504	A	20050907	CN 2003-815878	20030512
	JP 2005534633	T2	20051117	JP 2004-503004	20030512
	EP 1645274	A1	20060412	EP 2005-77643	20030512
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, SK				
	CA 2494518	AA	20040212	CA 2003-2494518	20030806
	WO 2004012739	A1	20040212	WO 2003-US24615	20030806
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

AU	2003258109	A1	20040223	AU 2003-258109	20030806
US	2004192700	A1	20040930	US 2003-635337	20030806
EP	1545530	A1	20050629	EP 2003-767250	20030806
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
CN	1688310	A	20051026	CN 2003-823620	20030806
JP	2006508909	T2	20060316	JP 2004-526061	20030806
US	2005288301	A1	20051229	US 2005-208248	20050819
US	2005288302	A1	20051229	US 2005-208299	20050819
PRAI US	2002-379510P	P	20020510		
US	2002-389093P	P	20020614		
US	2002-401672P	P	20020806		
US	2002-402749P	P	20020812		
US	2002-409860P	P	20020910		
US	2002-423338P	P	20021101		
US	2002-432961P	P	20021212		
US	2003-444812P	P	20030203		
US	2003-448062P	P	20030215		
EP	2003-750112	A3	20030512		
US	2003-436736	A3	20030512		
WO	2003-US14811	W	20030512		
WO	2003-US24615	W	20030806		

AB Provided are novel **crystalline** forms of **gatifloxacin** denominated forms A, B, C, D, E1, F, G, H, I, and J, and methods for their preparation Also provided are methods for making known **crystalline** forms of **gatifloxacin**, in particular forms omega and T2RP. Form A of gatifloxacin was prepared from a slurry in isopropanol.

IC ICM A61K031-496
ICS C07D401-04

CC 63-6 (Pharmaceuticals)
Section cross-reference(s): 75

ST **gatifloxacin crystal** form

IT **Crystal** morphology
(**crystalline** forms of **gatifloxacin**)

IT **112811-59-3**, **Gatifloxacin** 180200-66-2, **Gatifloxacin** sesquihydrate 404858-36-2, **Gatifloxacin** hemihydrate 614751-80-3, 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo-, hydrate 624736-92-1 624736-93-2 624736-94-3 624736-95-4
RL: FMU (Formation, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); FORM (Formation, nonpreparative); USES (Uses)
(**crystalline** forms of **gatifloxacin**)

IT 64-17-5, Ethanol, processes 67-56-1, Methanol, processes 67-63-0, Isopropanol, processes 67-64-1, Acetone, processes 71-36-3, 1-Butanol, processes 78-93-3, Mek, processes 118240-86-1, Methanol-water mixture
RL: PEP (Physical, engineering or chemical process); PYP (Physical process); PROC (Process)
(**crystalline** forms of **gatifloxacin**)

IT **112811-59-3**, **Gatifloxacin**
RL: FMU (Formation, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); FORM (Formation, nonpreparative); USES (Uses)
(**crystalline** forms of **gatifloxacin**)

RN 112811-59-3 HCAPLUS

CN 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-

(3-methyl-1-piperazinyl)-4-oxo- (9CI) (CA INDEX NAME)



L14 ANSWER 11 OF 16 HCAPLUS COPYRIGHT 2006 ACS on STN
AN 2003:836855 HCAPLUS
DN 139:328373
TI Anhydrous crystalline forms I and II of gatifloxacin
IN Reddy, Manne Satyanarayana; Raju, Chakilam Naga; Raju, Vetukuri Venkata Naga Kali Vara Prasada; Reddy, Ningam Srinivas; Kumar, Rapolu Rajesh
PA Reddy's Laboratories Limited, India; Cord, Janet I.
SO PCT Int. Appl., 25 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003086402	A1	20031023	WO 2003-US10708	20030407
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CA 2481308	AA	20031023	CA 2003-2481308	20030407
	AU 2003230831	A1	20031027	AU 2003-230831	20030407
	EP 1492535	A1	20050105	EP 2003-723931	20030407
	EP 1492535	B1	20051026		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
	AT 307583	E	20051115	AT 2003-723931	20030407
	US 2006142300	A1	20060629	US 2005-510892	20050908
PRAI	IN 2002-MA259	A	20020408		
	IN 2002-MA285	A	20020412		
	WO 2003-US10708	W	20030407		

GI



I

AB The present invention relates to the novel anhydrous crystalline forms I and II of gatifloxacin (I). The present invention also relates to methods of making the anhydrous Forms I and II of I, use of the forms and methods for preparing them.

IC ICM A61K031-496
ICS C07D215-56

CC 63-6 (Pharmaceuticals)
Section cross-reference(s): 28, 75

ST **gatifloxacin crystal** form

IT Crystal morphology
(anhydrous crystalline forms I and II of gatifloxacin)

IT **112811-59-3P, Gatifloxacin**
RL: PEP (Physical, engineering or chemical process); PRP (Properties); PYP (Physical process); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)
(anhydrous **crystalline** forms I and II of gatifloxacin)

IT 71-43-2, Benzene, processes 75-97-8 78-93-3, MEK, processes 108-10-1, MIBK 108-88-3, Toluene, processes 110-82-7, Cyclohexane, processes 1330-20-7, Xylene, processes
RL: PEP (Physical, engineering or chemical process); PYP (Physical process); PROC (Process)
(anhydrous crystalline forms I and II of gatifloxacin)

IT 109-07-9, 2-Methylpiperazine 112811-72-0, 3-Quinolinecarboxylic acid, 1-cyclopropyl-6,7-difluoro-1,4-dihydro-8-methoxy-4-oxo- 614751-80-3
RL: RCT (Reactant); RACT (Reactant or reagent)
(anhydrous crystalline forms I and II of gatifloxacin)

IT **112811-59-3P, Gatifloxacin**
RL: PEP (Physical, engineering or chemical process); PRP (Properties); PYP (Physical process); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)
(anhydrous **crystalline** forms I and II of gatifloxacin)

RN 112811-59-3 HCAPLUS

CN 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo- (9CI) (CA INDEX NAME)



RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L14 ANSWER 12 OF 16 HCAPLUS COPYRIGHT 2006 ACS on STN
AN 2003:5707 HCAPLUS
DN 138:61343
TI Pediatric formulation of gatifloxacin
IN Raghavan, Krishnaswamy S.; Ranadive, Sunanda A.; Bembenek, Kenneth S.; Benkerrour, Loutfy; Trognon, Veronique; Corrao, Richard G.; Esposito, Luigi
PA Bristol-Myers Squibb Company, USA
SO PCT Int. Appl., 16 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003000175	A2	20030103	WO 2002-US14596	20020510
	WO 2003000175	A3	20030410		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	US 2003028025	A1	20030206	US 2002-143487	20020509
	US 6589955	B2	20030708		
	CA 2450742	AA	20030103	CA 2002-2450742	20020510
	NZ 529611	A	20031219	NZ 2002-529611	20020510
	EP 1406572	A2	20040414	EP 2002-744141	20020510
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
	EE 200400018	A	20040415	EE 2004-18	20020510
	BR 2002009692	A	20040420	BR 2002-9692	20020510
	CN 1518449	A	20040804	CN 2002-812409	20020510
	JP 2005500306	T2	20050106	JP 2003-506621	20020510
	TW 224004	B1	20041121	TW 2002-91113074	20020614
	ZA 2003009173	A	20050225	ZA 2003-9173	20031125
	BG 108444	A	20040831	BG 2003-108444	20031212
PRAI	US 2001-299625P	P	20010620		
	WO 2002-US14596	W	20020510		

AB A taste-masked formulation of the quinolone antibacterial gatifloxacin for pediatric uses is described. A **crystalline** co-precipitate of **gatifloxacin** and one or both of stearic acid and palmitic acid in a narrow weight ratio has been found to effectively mask the bitter taste of gatifloxacin. The taste of gatifloxacin is effectively masked in the mouth and in aqueous suspension through a full dosage cycle, typically 14 days. **Gatifloxacin** in the subject **crystalline** co-ppts. has been found to be readily available for absorption from the stomach.
IC ICM A61K
CC 63-6 (Pharmaceuticals)
ST gatifloxacin palmitate stearate ppt taste suspension child
IT Development, mammalian postnatal
(child; **gatifloxacin crystalline** co-ppts. with stearic

acid and/or palmitic acid for taste-masked pediatric formulations)
IT Drug delivery systems
(suspensions, oral; **gatifloxacin crystalline** co-ppts. with stearic acid and/or palmitic acid for taste-masked pediatric formulations)
IT 57-10-3, Palmitic acid, biological studies 57-11-4, Stearic acid, biological studies
RL: MOA (Modifier or additive use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(**gatifloxacin crystalline** co-ppts. with stearic acid and/or palmitic acid for taste-masked pediatric formulations)
IT **112811-59-3, Gatifloxacin**
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(**gatifloxacin crystalline** co-ppts. with stearic acid and/or palmitic acid for taste-masked pediatric formulations)
IT **112811-59-3, Gatifloxacin**
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(**gatifloxacin crystalline** co-ppts. with stearic acid and/or palmitic acid for taste-masked pediatric formulations)
RN 112811-59-3 HCAPLUS
CN 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo- (9CI) (CA INDEX NAME)



L14 ANSWER 13 OF 16 HCAPLUS COPYRIGHT 2006 ACS on STN
AN 2002:831071 HCAPLUS
DN 139:78466
TI Influence of clarithromycin on biofilm of Pseudomonas aeruginosa
AU Wang, Rui; Pei, Fei; Chai, Dong; Zhu, Man; Fang, Yi; Wang, Zhongxiao; Liu, Guiyang
CS The General Hospital of PLA, Beijing, 100853, Peop. Rep. China
SO Zhongguo Kangshengsu Zazhi (2002), 27(5), 293-297
CODEN: ZKZAEY; ISSN: 1001-8689
PB Zhongguo Kangshengsu Zazhishe
DT Journal
LA Chinese
AB The influence of clarithromycin on Pseudomonas aeruginosa biofilm was studied. The adherence of mucoid Pseudomonas aeruginosa was measured by **crystal** violet staining method, the **gatifloxacin** penetration concentration was determined by HPLC, and the effect of clarithromycin on Pseudomonas aeruginosa biofilm was determined by carbazole/ethanol method. The stable biofilm could be formed in 7 days. When 8 x MIC gatifloxacin was combined with 1/4 MIC clarithromycin, the adherence of mucoid Pseudomonas aeruginosa was significantly decreased, and the optical d. was decreased from (0.126 ± 0.011) to (0.114 ± 0.010). The penetrated concns. of gatifloxacin through biofilm were increased from (1.210 ± 0.091), (2.911 ± 0.112), and (5.911 ± 0.213) to (1.752 ± 0.122), (3.908

± 0.154), and (7.898 ± 0.321) μg mL-1 in three dosages of 4, 8, and 16 μg mL-1, resp., and the concentration of alginate in biofilm was decreased from (10.07 ± 0.55) to (2.34 ± 0.21), (4.91 ± 0.16), (7.22 ± 0.36), and (8.82 ± 0.50) μg per 108 CFU in 80, 40, 20, and 10 μg mL-1 clarithromycin groups, resp. Clarithromycin could reduce the adherence of Pseudomonas aeruginosa, enhance the penetration ability of gatifloxacin, and inhibit the alginate synthesis of Pseudomonas aeruginosa.

CC 1-5 (Pharmacology)

ST clarithromycin Pseudomonas biofilm gatifloxacin alginate

IT Adhesion, biological
Biofilms (microbial)
Pseudomonas aeruginosa
(influence of clarithromycin on biofilm of Pseudomonas aeruginosa)

IT 81103-11-9, Clarithromycin
RL: BUU (Biological use, unclassified); DMA (Drug mechanism of action); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(influence of clarithromycin on biofilm of Pseudomonas aeruginosa)

IT **112811-59-3**, Gatifloxacin
RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(penetration of biofilm; influence of clarithromycin on biofilm of Pseudomonas aeruginosa)

IT 9005-32-7, Alginic acid
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(response; influence of clarithromycin on biofilm of Pseudomonas aeruginosa)

IT **112811-59-3**, Gatifloxacin
RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(penetration of biofilm; influence of clarithromycin on biofilm of Pseudomonas aeruginosa)

RN 112811-59-3 HCAPLUS

CN 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo- (9CI) (CA INDEX NAME)



L14 ANSWER 14 OF 16 HCAPLUS COPYRIGHT 2006 ACS on STN

AN 2002:220371 HCAPLUS

DN 136:268136

TI Preparation of gatifloxacin pentahydrate

IN Raghavan, Krishnaswamy S.; Ranadive, Sunanda A.; Gougoutas, Jack Z.; Dimarco, John D.; Parker, William L.; Davidovich, Martha; Neuman, Ann

PA Bristol-Myers Squibb Company, USA

SO PCT Int. Appl., 20 pp.
CODEN: PIXXD2

DT Patent

LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002022126	A1	20020321	WO 2001-US26120	20010821
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	US 2002052379	A1	20020502	US 2001-932045	20010817
	US 6413969	B2	20020702		
	CA 2422616	AA	20020321	CA 2001-2422616	20010821
	AU 2001086592	A5	20020326	AU 2001-86592	20010821
	EP 1326612	A1	20030716	EP 2001-966049	20010821
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
	JP 2004508403	T2	20040318	JP 2002-526377	20010821
	BR 2001013866	A	20040706	BR 2001-13866	20010821
	CN 1511035	A	20040707	CN 2001-818780	20010821
PRAI	US 2000-232293P	P	20000913		
	WO 2001-US26120	W	20010821		

AB **Crystalline gatifloxacin** pentahydrate (I) in a highly homogeneous form with respect to other crystalline forms is disclosed. Thus, I was prepared by suspending gatifloxacin hemihydrate in water, filtered, and the product was dried for 16 h. Thus, a tablet composition contained I 0.428, microcryst. cellulose 0.138, sodium starch glycolate 0.024, and Mg stearate 0.09 g/tablet.

IC ICM A61K031-496
ICS C07D401-00

CC 63-6 (Pharmaceuticals)

ST gatifloxacin pentahydrate pharmaceutical prepn

IT Drug delivery systems
(oral; preparation of gatifloxacin pentahydrate)

IT Drug delivery systems
(parenterals; preparation of gatifloxacin pentahydrate)

IT **Crystal** morphology
(preparation of **gatifloxacin** pentahydrate)

IT Drug delivery systems
(solids, oral; preparation of gatifloxacin pentahydrate)

IT Drug delivery systems
(suspensions, oral; preparation of gatifloxacin pentahydrate)

IT Drug delivery systems
(suspensions; preparation of gatifloxacin pentahydrate)

IT Drug delivery systems
(tablets; preparation of gatifloxacin pentahydrate)

IT 404858-35-1P, Gatifloxacin pentahydrate
RL: IMF (Industrial manufacture); PRP (Properties); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of gatifloxacin pentahydrate)

IT **112811-59-3**, Gatifloxacin 180200-66-2, Gatifloxacin sesquihydrate 404858-36-2, Gatifloxacin hemihydrate
RL: RCT (Reactant); THU (Therapeutic use); BIOL (Biological study); RACT (Reactant or reagent); USES (Uses)
(preparation of gatifloxacin pentahydrate)

IT **112811-59-3**, Gatifloxacin
RL: RCT (Reactant); THU (Therapeutic use); BIOL (Biological study); RACT (Reactant or reagent); USES (Uses)
(preparation of gatifloxacin pentahydrate)
RN 112811-59-3 HCAPLUS
CN 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo- (9CI) (CA INDEX NAME)



RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L14 ANSWER 15 OF 16 HCAPLUS COPYRIGHT 2006 ACS on STN
AN 2002:180387 HCAPLUS
DN 137:60163
TI Influence of anti-alginate serum on adherence and penetration of mucoid Pseudomonas aeruginosa biofilm
AU Pei, Fei; Wang, Rui; Chai, Dong; Fang, Yi; Liu, Guiyang; Zhu, Man; Li, Cunfu; Di, Min
CS Department of Clinical Pharmacology of General Hospital of PLA, Beijing, 100853, Peop. Rep. China
SO Zhongguo Linchuang Yaolixue Zazhi (2001), 17(6), 423-426
CODEN: ZLYZE9; ISSN: 1001-6821
PB Beijing Yike Daxue, Linchuang Yaoli Yanjiuso
DT Journal
LA Chinese
AB The influence of anti-alginate serum on the adherence and the penetration of mucoid Pseudomonas aeruginosa biofilm were assessed. The adherence of mucoid Pseudomonas aeruginosa was determined by **crystal** violet stained method; the **gatifloxacin** penetration concentration was determined by HPLC. When 1:1 dilution anti-alginate serum combined with 8 x MIC of gatifloxacin, the adherence of mucoid Pseudomonas aeruginosa decreased significantly, the optical d. of silicon slides at 540 nm decreased from 0.130±0.010, 0.129±0.015 and 0.126±0.011 to 0.120±0.010, 0.117±0.015 and 0.1140±0.010. The penetrated concns. of gatifloxacin through biofilm were increased from 1.210±0.091, 2.911±0.112 and 5.911±0.213 to 1.752±0.122, 3.908±0.154 and 7.898±0.321. The anti-alginate serum could reduce the adherence of Pseudomonas aeruginosa and enhance the penetration ability of gatifloxacin.
CC 10-6 (Microbial, Algal, and Fungal Biochemistry)
ST anti alginate serum biofilm adherence penetration Pseudomonas; mucoid Pseudomonas aeruginosa
IT Adhesion, biological
Antiserums
Biofilms (microbial)
Pseudomonas aeruginosa
(influence of anti-alginate serum on adherence and penetration of mucoid Pseudomonas aeruginosa biofilm)

IT 9005-32-7, Alginic acid **112811-59-3**, Gatifloxacin
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(influence of anti-alginate serum on adherence and penetration of mucoid Pseudomonas aeruginosa biofilm)
IT **112811-59-3**, Gatifloxacin
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(influence of anti-alginate serum on adherence and penetration of mucoid Pseudomonas aeruginosa biofilm)
RN 112811-59-3 HCAPLUS
CN 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo- (9CI) (CA INDEX NAME)



L14 ANSWER 16 OF 16 HCAPLUS COPYRIGHT 2006 ACS on STN
AN 2000:144737 HCAPLUS
DN 132:185458
TI Aqueous liquid preparations of gatifloxacin
IN Yasueda, Shinichi; Inada, Katsuhiro
PA Senju Pharmaceutical Co., Ltd., Japan; Kyorin Pharmaceutical Co., Ltd.
SO PCT Int. Appl., 17 pp.
CODEN: PIXXD2
DT Patent
LA Japanese
FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 2000010570	A1	20000302	WO 1999-JP4483	19990820
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
CA 2307632	AA	20000302	CA 1999-2307632	19990820
AU 9953026	A1	20000314	AU 1999-53026	19990820
AU 761040	B2	20030529		
EP 1025846	A1	20000809	EP 1999-938550	19990820
EP 1025846	B1	20060712		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY				
BR 9906735	A	20000815	BR 1999-6735	19990820
NZ 504017	A	20010928	NZ 1999-504017	19990820
TW 537895	B	20030621	TW 1999-88114247	19990820
CN 1133432	B	20040107	CN 1999-801408	19990820
AT 332692	E	20060815	AT 1999-938550	19990820
US 6333045	B1	20011225	US 2000-529882	20000421

PRAI JP 1998-235432 A 19980821
WO 1999-JP4483 W 19990820

AB This invention relates to aqueous prepns. containing gatifloxacin [(±)-1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo-quinolinecarboxylic acid] or its salts and sodium edetate. Also disclosed are a method for enhancing the corneal permeability of **gatifloxacin**, a method for preventing **crystallization** of **gatifloxacin** and a method for preventing coloration of gatifloxacin each by blending gatifloxacin or its salt with sodium edetate. An aqueous solution for eye drops, ear drops, and nasal drops, contained gatifloxacin 0.5, Na edetate 0.1, NaCl 0.9, HCl/NaOH q.s. to pH 7, and sterilized water to 100 mL.

IC ICM A61K031-495
ICS A61K009-08; A61K047-18; C07D401-04

CC 63-6 (Pharmaceuticals)

ST gatifloxacin edetate stabilizer aq soln

IT Drug delivery systems
(solns., ear; stabilized aqueous prepns. containing gatifloxacin and edetate)

IT Drug delivery systems
(solns., nasal; stabilized aqueous prepns. containing gatifloxacin and edetate)

IT Drug delivery systems
(solns., ophthalmic; stabilized aqueous prepns. containing gatifloxacin and edetate)

IT Discoloration prevention agents
(stabilized aqueous prepns. containing gatifloxacin and edetate)

IT 64-02-8, Sodium edetate **112811-59-3**, Gatifloxacin
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(stabilized aqueous prepns. containing gatifloxacin and edetate)

IT **112811-59-3**, Gatifloxacin
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(stabilized aqueous prepns. containing gatifloxacin and edetate)

RN 112811-59-3 HCAPLUS

CN 3-Quinolinecarboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-4-oxo- (9CI) (CA INDEX NAME)



RE.CNT 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

=>